

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/IB2006/051199

International filing date: 18 April 2006 (18.04.2006)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/672,900  
Filing date: 18 April 2005 (18.04.2005)


Date of receipt at the International Bureau: 12 May 2006 (12.05.2006)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

PA 7008618



IB/06/54139

12 May 2006

# THE UNITED STATES OF AMERICA

**TO ALL TO WHOM THESE PRESENTS SHALL COME:  
UNITED STATES DEPARTMENT OF COMMERCE**

**United States Patent and Trademark Office**

**April 21, 2006**

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM  
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK  
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT  
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A  
FILING DATE UNDER 35 USC 111.**

**APPLICATION NUMBER: 60/672,900**

**FILING DATE: April 18, 2005**

**REC'D 12 MAY 2006**

**WIPO PCT RO/IB**

**THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY  
APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS  
CONVENTION, IS US60/672,900**

**By Authority of the  
Under Secretary of Commerce for Intellectual Property  
and Director of the United States Patent and Trademark Office**



*W. Montgomery*  
**W. MONTGOMERY**  
**Certifying Officer**



20427 U.S.

EV600680256US

PTO/SB/16 (12-04)  
Approved for use through 07/31/2006. OMB 0651-0032  
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE  
This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.  
**PROVISIONAL APPLICATION FOR PATENT COVER SHEET**  
This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).  
Express Mail Label No. EV 600680256 US

INVENTOR(S)		
Given Name (first and middle (if any))	Family Name or Surname	Residence (City and either State or Foreign Country)
Filippo	Giancotti	New York, NY
Additional inventors are being named on the _____ separately numbered sheets attached hereto		
TITLE OF THE INVENTION (500 characters max)		
Inhibition of tumorigenesis by inhibition of a6b4 integrin		
Direct all correspondence to: CORRESPONDENCE ADDRESS		
<input checked="" type="checkbox"/> The address corresponding to Customer Number: 021121		
OR		
<input type="checkbox"/> Firm or Individual Name Marina T. Larson, Ph.D		
Address		
City	State	Zip
Country	Telephone	Fax
ENCLOSED APPLICATION PARTS (check all that apply)		
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76		
<input checked="" type="checkbox"/> Specification. Number of Pages 17		
<input checked="" type="checkbox"/> Drawing(s) Number of Sheets 10		
<input type="checkbox"/> CD(s), Number of CDs _____		
<input type="checkbox"/> Other (specify) _____		
Application Size Fee: If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$ 250 (\$ 125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).		
METHOD OF PAYMENT OF FILING FEES AND APPLICATION SIZE FEE FOR THIS PROVISIONAL APPLICATION FOR PATENT		
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.		
TOTAL FEE AMOUNT \$		
<input type="checkbox"/> A check or money order is enclosed to cover the filing fee and application size fee (if applicable). 100.00		
<input checked="" type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.		
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fee and application size fee (if applicable) or credit any overpayment to Deposit Account Number. 15-0610. A duplicative copy of this form is enclosed for fee processing.		
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.		
<input type="checkbox"/> No.		
<input checked="" type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: NIH Merit Award R13 CA58976		

SIGNATURE Marina T. LarsonDate April 18, 2005TYPED or PRINTED NAME Marina T. Larson, Ph.DREGISTRATION NO. 32038  
(if appropriate)TELEPHONE 970-468-6600Docket Number: MSK-P-082-PV**USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT**

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

## **Inhibition of tumorigenesis by inhibition of $\alpha 6 \beta 4$ integrin**

### **Background of Invention**

This application relates to a method of inhibiting tumorigenesis, particularly in the case of tyrosine kinase-related cancers such as breast and prostate cancer, through the inhibition of  $\alpha 6 \beta 4$  integrin using a therapeutic agent that targets the  $\beta 4$  portion of the integrin. In this application, the nomenclature  $\alpha 6 \beta 4$  refers to the  $\alpha 6$ - $\beta 4$  integrin. Similar nomenclature with arabic or roman numerals is used for other integrins.

Integrins are a class of cellular transmembrane receptors known to bind to extracellular matrix proteins, and therefore they mediate cell-cell and cell-extracellular matrix interactions, referred generally to as cell adhesion events. The integrins connect the extracellular matrix to the intracellular cytoskeleton and cooperate with Receptor Protein Tyrosine Kinases (RPTKs) to regulate cell fate (Giancotti and Ruoslahti, 1999; Hynes, 2003; Miranti and Brugge, 2002). Depending on the integrins they express and the matrix they attach to, normal cells proliferate or undergo growth arrest, migrate or remain stationary, and live or undergo apoptotic death. These effects imply that the integrins impart a stringent control to the action of RPTKs, determining the nature and direction of the cell's response to growth factors and cytokines (Giancotti and Tarone, 2003). Despite considerable amounts of cell biological data, genetic evidence of the significance of integrin signaling remains scarce. In particular, it has been difficult to separate the adhesive and signaling function of individual integrins in any model system analyzed to date.

The integrin receptors constitute a family of proteins with shared structural characteristics of noncovalent heterodimeric glycoprotein complexes formed of  $\alpha$  and  $\beta$  subunits. There are eight known  $\beta$  subunits and fourteen known  $\alpha$  subunits, which associate in various combinations to form twenty five receptors with different ligand specificities. The ligands for several of the integrins are adhesive extracellular matrix (ECM) proteins such as fibronectin, vitronectin, collagens and laminin. It has been reported that the  $\alpha 5 \beta 1$  integrin, a fibronectin receptor, and the  $\alpha$  integrins  $\alpha 5 \beta 3$  and  $\alpha 5 \beta 5$ , which bind to several RGD-containing matrix proteins, promote angiogenesis. Hynes et al. (2002). This property has been considered as a basis for using inhibitors of such integrins as inhibitors of angiogenesis. See US Patents Nos. 5,981,478; 5,766,591; 6,358,970; and 6,645,991. However, while genetic experiments in mice have confirmed the role of  $\alpha 5 \beta 1$  integrin in angiogenesis, they have not confirmed a role for the  $\alpha$  integrins, thus calling into question the efficacy of anti-angiogenic therapy based on the latter group. Anti-angiogenic therapy based on

inhibition of  $\alpha 5 \beta 1$  integrin is problematic because of toxicity arising as a result of the critical involvement of this integrin adhesion of several cell types.

The  $\alpha 6 \beta 4$  integrin is a laminin-5 receptor expressed by epithelial cells, Schwann cells, and endothelial cells and has several distinguishing features. The cytoplasmic domain of  $\beta 4$  is unusually long (ca. 1000 amino acids) and displays no homology to the short cytoplasmic tails of other  $\beta$  subunits. Upon  $\alpha 6 \beta 4$  binding to matrix, the unique cytoplasmic domain of  $\beta 4$  is phosphorylated on multiple tyrosines by a Src Family kinase (SFK) and interacts directly with the signaling adaptor protein Shc, causing activation of the Ras to ERK cascade (Dans et al., 2001; Gagnoux-Palacios et al., 2003; Mainiero et al., 1995). In addition, the  $\beta 4$  tail mediates activation of PI-3K and Rac (Shaw, 2001; Shaw et al., 1997). Upon dephosphorylation, the cytoplasmic domain of  $\beta 4$  associates with the keratin cytoskeleton, causing assembly of hemidesmosomes and, hence, strengthening adhesion to laminin-5-containing basement membranes (Dans et al., 2001; Murgia et al., 1998; Spinardi et al., 1993). In contrast, the other integrins activate FAK/SFK signaling at focal adhesions (Geiger et al., 2001; Schlaepfer and Hunter, 1998) and, although some of them also recruit Shc, they do so by a distinct, indirect mechanism (Wary et al., 1998).

The pattern of expression of  $\alpha 6 \beta 4$  in the skin is consistent with a role for  $\alpha 6 \beta 4$  signaling in the control of epithelial proliferation. In normal epidermis, the expression of  $\alpha 6 \beta 4$  is restricted to the basal cell layer, which comprises the rapidly dividing transit-amplifying cells (Borradori and Sonnenberg, 1999; Fuchs et al., 1997), while in skin diseases characterized by suprabasal proliferation, such as squamous carcinoma and psoriasis,  $\alpha 6 \beta 4$  extends to the suprabasal layers (Pellegrini et al., 1992). In addition, ligation of  $\alpha 6 \beta 4$  promotes progression through G1 and entry in S phase in keratinocytes treated with EGF, whereas ligation of  $\alpha 2 \beta 1$  does not exert this effect (Mainiero et al., 1997).

Tumor biology studies have suggested a function for  $\alpha 6 \beta 4$  signaling in tumor invasion. Many invasive carcinomas display elevated levels of  $\alpha 6 \beta 4$  (Mercurio and Rabinovitz, 2001). Introduction of  $\alpha 6 \beta 4$  in breast and colon carcinoma cells that have lost its expression activates PI-3K to Rac signaling and increases invasive ability in vitro (Shaw et al., 1997). In addition, it has been proposed that the  $\beta 4$  tail functions as an essential adapter and amplifier of pro-invasive signals elicited by activated Met in cells undergoing Met-induced oncogenesis (Trusolino et al., 2001). Finally, introduction of a dominant negative form of  $\beta 4$  impairs the survival of breast carcinoma cells, and this effect has been linked to the ability of mutant  $\beta 4$  to interfere with the assembly of hemidesmosomes and the establishment of a partially polarized phenotype (Weaver et al., 2002). Collectively, these results suggest the possibility that  $\alpha 6 \beta 4$  promotes cell migration and invasion and confers resistance to apoptosis in carcinoma cells.

Commonly assigned US Provisional Application No. 60/481,696, filed November 22, 2003, and PCT application PCT/US2004/039189, which are incorporated herein by reference, described the use of  $\alpha 6 \beta 4$  integrin in controlling pathological neogenesis. While this angiogenesis can occur in tumors, controlling angiogenesis is not the same as controlling or inhibiting tumorigenesis and tumor progression.

It has been shown previously that coexpression of  $\alpha 6 \beta 4$  and laminin and amplification of ErbB-2 correlate with a poor prognosis in breast cancer patients (Slamon et al., 1987; Tagliabue et al., 1998). Although a significant fraction of human breast cancers show reduced expression of  $\alpha 6 \beta 4$ , a significant fraction of metastatic lymph nodes stain positive for  $\alpha 6 \beta 4$  (Natali et al., 1992). Further, there is evidence that introduction of  $\alpha 6 \beta 4$  increases the invasive ability of MDA-MB-435 breast carcinoma cells in in vitro assays (Shaw et al., 1997) and that the ability of  $\beta 4$ -transfected MDA-MB-435 cells to metastasize to lung upon injection into a mouse tail vein requires the binding of  $\alpha 6 \beta 4$  on cancer cells to Lu-ECAM-1 on lung endothelial cells. (Abdel-DGhany et al., 2001). Pathological studies have shown that the expression of  $\alpha 6 \beta 4$  declines during prostate cancer progression (Cress et al., Cancer Metastasis Rev. 14:219-228, 1995). However, the integrin is still expressed at significant levels in Prostate Intraepithelial Neoplasia (PIN), and it may play a role at this stage of tumor progression. Notwithstanding these findings, the role, if any, of  $\alpha 6 \beta 4$  in tumor progression is not understood in the art. For example, whether the observed variations in expression levels are cause or effect, whether reduction changes in  $\alpha 6 \beta 4$  have any actual impact of tumor growth or invasiveness in vivo, and how  $\alpha 6 \beta 4$  interacts with other signaling moieties are not known.

#### Summary of Invention

Following an investigation of the role of  $\alpha 6 \beta 4$  in mice engineered to develop mammary tumors on expression of an activated version of ErbB-2 and on mice engineered to develop prostate cancer on expression of the SV-40 T Antigen, we have determined that  $\alpha 6 \beta 4$  integrin signaling is necessary for the progression of breast and prostate cancer. This finding is also applicable to other tumor types that express  $\alpha 6 \beta 4$ , such as thyroid cancer, squamous carcinoma of the skin, cervix, and upper gastrointestinal tract, pancreatic cancer, colon cancer (Mercurio AM, Rabinovitz I. Towards a mechanistic understanding of tumor invasion--lessons from the  $\alpha 6 \beta 4$  integrin. Semin Cancer Biol. 2001 Apr;11(2):129-41). Thus, the present invention provides methods for the inhibition of tumorigenesis in tumors of this type using inhibitors of  $\alpha 6 \beta 4$  integrin that target  $\beta 4$ . In accordance with the method of the invention, an individual in whom tumorigenesis is to be inhibited is exposed to a therapeutic agent effective to reduce the amount of active  $\alpha 6 \beta 4$  integrin in the individual, at least at locations relevant to tumorigenesis. In one embodiment of the invention, the individual is a human patient. The therapeutic agent may be an antibody or a small molecule, for example a

laminin-5 analog, which binds to  $\alpha 6 \beta 4$  integrin and inhibits its normal function. The therapeutic agent may also be a chemical species that interferes with the production of  $\beta 4$ , including for example an antisense or RNAi species. The therapeutic agent is administered to the tissue or patient in a therapeutically effective amount. The therapeutic agent may be used as a single agent or in combination with other therapies, especially those directed toward suppressing the activity of RPTKs known to cooperate with  $\alpha 6 \beta 4$ , including but not limited to ErbB2, EGF-R, Met, and Ron.

#### **Brief Description of the Drawings**

Fig. 1 shows a breeding strategy for introduction of MMTV-Neu<sup>Ndl</sup>-YD transgene into both wild-type and  $\beta 4$ -1355T mice.

Fig. 2 shows the extent of tumor free survival in  $\beta 4$  mutant and wild-type breed in accordance with the scheme in Fig. 1.

Fig. 3 shows the number of individual mammary tumors in individual mice.

Fig. 4 shows the growth of mammary tumors in wild-type and  $\beta 4$ -mutant mice.

Fig. 5 shows the difference in histological progression in mammary tumors in wild-type and  $\beta 4$ -mutant mice.

Fig. 6 shows a breeding strategy for introduction of TRAMP into both wild-type and  $\beta 4$ -1355T mice.

Fig. 7 shows results of an MRI analysis indicative of tumor growth in  $\beta 4$ -mutant and  $\beta 4$ -wild-type TRAMP mice.

Fig. 8 shows survival of  $\beta 4$ -mutant and  $\beta 4$ -wild-type TRAMP mice.

Fig. 9 shows the sensitization upon loss of  $\beta 4$  signaling when MMTV-Neu (YD) mice bearing mammary tumors were treated with Iressa or vehicle (0.1% Tween-80).

Fig. 10 shows reduction in tumor volume when MMTV-Neu (YD) mice bearing mammary tumors were treated with Iressa.

#### **Detailed Description**

As used in this application, the term "tumorigenesis" refers to initiation of primary or metastatic tumor growth, and the promotion of invasive growth.

As used in this application, the term "inhibition" refers to a reduction of the event or activity inhibited to an extent sufficient to produce an observable result. Complete elimination of the event or activity is not required.

As used in this application, the term "amount of active  $\alpha 6 \beta 4$  integrin" refers to the observable tumorigenesis-promoting activity resulting from  $\alpha 6 \beta 4$  integrin present in a tissue. Reductions in the amount of the active  $\alpha 6 \beta 4$  integrin can result from a reduction in the amount of  $\alpha 6 \beta 4$  integrin, i.e., effectively a reduction in concentration; a reduction in the capacity of individual molecules of  $\alpha 6 \beta 4$  integrin to promote tumorigenesis, i.e., effectively a change in the quality of the integrin, or combinations thereof. The first type of reduction will most commonly be achieved by limiting the production of  $\alpha 6 \beta 4$  integrin, for example using an antisense oligonucleotide or RNAi techniques, although it could also be achieved by accelerating the decomposition of  $\alpha 6 \beta 4$  integrin. The second type of reduction is most readily achieved through physical binding of the integrin with a ligand that competes with the normal ligand for binding to the receptor.

As used in this application, the terms "treatment" or "treating" refer to the application of a therapeutic agent to achieve a reduction in the amount of active  $\alpha 6 \beta 4$  integrin so as to produce a benefit to a patient being treated. Such a benefit need not be a complete or permanent cure, but may be only a lessening of the rate at which tumorigenesis is occurring, thereby delaying progression of a disease condition.

As used in this application, the term "administration" refers to any means by which a therapeutic agent can be delivered to a tissue, including without limitation oral, nasal and transdermal administration and injection, for example subcutaneous, subdermal, intramuscular, intravenous, intrathecal or peritoneal injection. For treatment of eye-associated tumorigenesis, direct injection to the eye may be used. The therapeutic agent of the invention can be used in combination with other agents used in the treatment of cancer. In particular, the therapeutic agent of the invention is suitably used in combination with kinase inhibitors such as Iressa. Use in combination entails the administration of two or more agents in a time course where the effects of at least one of the agents is improved as a result of the use of the other. Two agents need not be administered at the same time to be considered use in combination, and may be used in any order.

The effective amount of a therapeutic agent to be administered varies depending on the nature of the therapeutic agent, and will frequently reflect a balancing of therapeutic benefits and side effects. However, the determination of specific amounts for a given therapeutic is routine and within the skill in the art.



Therapeutic agents useful in the present invention may be antibodies, aptamers or small molecules that bind to  $\alpha 6 \beta 4$  integrin to produce a reduction in activity. Examples include small molecules which block  $\beta 4$  signalling by binding to  $\beta 4$ , and have specific functions such as inhibiting nuclear translocation of NF- $\kappa$ B. Where an antibody therapeutic agent is used, it may be administered in the form of the antibody, or formed in situ by expression of a nucleic acid sequence encoding an  $\alpha 6 \beta 4$  integrin-specific antibody. Such antibodies may be monoclonal, polyclonal, or modified constructs, for example single chain Fv constructs, targeting  $\alpha 6 \beta 4$  integrin. Binding sites may be on the alpha chain, the beta chain or both chains of the  $\alpha 6 \beta 4$  integrin. Non-antibody binding proteins could also be employed. For example, human Integrin- $\beta 4$  binding protein is known and has the sequence:

MAVRASFENNCEIGCFAKLTNTYCLVAIGGSENFYSVFEGELSDTIPVVHASIAGCRIIGRMCVGNRHG  
LLVPNNTTDQELQHIRNSLPDVTQIRRV EERLSALGNVTTCNDYVALVHPDL DRETEEILADV LKVEVF  
RQT VADQVLVGSYCVFSNQGLVHPKTSIEDQDELSSLLQVPLVAGTVNRGSEVIAAGMVVNDWCAF  
CGLDTTSTELSVVESVFKLNEAQPSTIATSMRDSLIDSLT(NM\_002212). (Seq. ID. No. 1).

The therapeutic agent may also be a nucleic acid that results in a reduction in the amount of active  $\alpha 6 \beta 4$  integrin, for example an antisense oligonucleotide or an RNA molecule that works by an RNAi mechanism. The nucleic acid may target, via a sequence specific mechanism, the alpha chain or the beta chain. The coding sequence of the beta 4 chain of human integrin is known from NM\_000213 to be as follows:

```

1 atggcagggc cagccccag cccatgggcc aggctgctcc tggcagcctt gatcagcgtc
61 agcctctctg ggaccttggc aaaccgctgc aagaaggccc cagtgaagag ctgcacggag
121 tgtgtccgtg tggataagga ctgcgcctac tgcacagacg agatgttcag ggaccggcgc
181 tgcaacaccc aggcggagct gctggccgcg ggctgccagc gggagagcat cgtggtcag
241 gagagcagct tccaaatcac agaggagacc cagattgaca ccaccctgcg gcgcagccag
301 atgtccccc aaggcctgcg ggtccgtctg cggcccgtg aggagcggca tttgagctg
361 gaggtgtttg agccactgga gagccccgtg gacctgtaca tcctcatgga cttctcaac
421 tccatgtccg atgatctgga caacctcaag aagatggggc agaactggc tcgggtcctg
481 agccagctca ccagcgacta cactattgga ttggcaagt ttgtggacaa agtcagcgtc
541 ccgcagacgg acatgaggcc tgagaagctg aaggagccct ggccaacag tgaccccccc
601 ttctcttca agaacgtcat cagcctgaca gaagatgtgg atgagttccg gaataaactg
661 cagggagagc ggaatctcag caacctggat gctcctgagg gcggcttcga tgccatcctg
721 cagacagctg tgtgcacgag ggacattggc tggcgcccgg acagcaccca cctgctggtc
781 ttctccaccg agtcagcctt ccactatgag gctgatggcg ccaacgtgct ggctggcatc
841 atgagccgca acgatgaacg gtgccacctg gacaccacgg gcacctacac ccagtacagg
901 acacaggact acccgctcgt gccaccctg gtgcgcctgc tcgccaagca caacatcatc
961 cccatctttg ctgtcaccaa ctactctat agctactacg agaagcttca cacctatttc
1021 cctgtctcct cactgggggt gctgcaggag gactcgtcca acatcgtgga gctgctggag

```

1081 gaggccttca atcggatccg ctccaacctg gacatccggg ccctagacag cccccgaggc  
 1141 ctccggacag aggtcacctc caagatgtc cagaagacga ggactgggtc cttcacatc  
 1201 cggcgggggg aagtgggtat ataccaggtg cagctgcggg cccttgagca cgtggatggg  
 1261 acgcacgtgt gccagctgcc ggaggaccag aagggcaaca tccatctgaa accttccttc  
 1321 tccgacggcc tcaagatgga cgcgggcac atctgtgatg tgtgcacctg cgagctgcaa  
 1381 aaagaggtgc ggtcagctcg ctgcagcttc aacggagact tcgtgtgcgg acagtgtgtg  
 1441 tgcagcgagg gctggagtgg ccagacctgc aactgtcca ccggctctct gagtgcatt  
 1501 cagccctgcc tgcgggaggg cgaggacaag ccgtgtccg gccgtgggga gtgccagtgc  
 1561 gggcactgtg tgtgtacgg cgaaggccgc tacgagggtc agttctgcga gtatgacaac  
 1621 ttccagtgc cccgcacttc cgggttctg tgcaatgacc gaggacgtg ctccatgggc  
 1681 cagtgtgtg gtgagcctgg ttggacaggc ccaagctgtg actgtccct cagcaatgcc  
 1741 acctgcatcg acagcaatgg gggcatctgt aatggacgtg gccactgtga gtgtggccg  
 1801 tgccactgcc accagcagtc gctctacag gacaccatct gcgagatcaa ctactcggc  
 1861 atccaccgg gcctctgcga ggacctacg tctgcgtgc agtgccaggc gtggggcacc  
 1921 ggcgagaaga aggggcgcac gtgtgaggaa tgcaactca aggtcaagat ggtggacgag  
 1981 cttagagag ccgaggaggt ggtggtgcgc tgcctctcc gggacgagga tgacgactgc  
 2041 acctacagct acaccatgga aggtgacggc gccctgggc ccaacagcac tgcctgggtg  
 2101 cacaagaaga aggactgcc tccgggctcc ttctggggc tcacccct gctcctctc  
 2161 ctctgccgc tctggccct gctactgtg ctatgtgga agtactgtc ctgctgcaag  
 2221 gcctgctgg cacttctcc gtgctgcaac cgaggtcaca tgggggctt taaggagac  
 2281 cactacatgc tgcgggagaa cctgatggc tctgacct tggacagcc catgctgcg  
 2341 agcgggaacc tcaagggccg tgacgtggtc cgctggaagg tcaccaaca catgcagcgg  
 2401 cctggctttg ccactcatgc cgccagcatc aacccacag agctgggtcc ctacgggctg  
 2461 tccttgccc tggccgcct ttgcaccgag aacctgtga agcctgacac tcgggagtg  
 2521 gccagctgc gccaggaggt ggaggagaac ctgaacgagg tctacaggca gatctccgg  
 2581 gtacacaagc tccagcagac caagttccg cagcagccca atgccgggaa aaagcaagac  
 2641 cacaccattg tggacacagt gctgatggc cccgctcg ccaagccggc cctgtgaag  
 2701 cttacagaga agcaggtgga acagagggcc ttccagacc tcaaggtggc cccggctac  
 2761 tacacctca ctgcagacca ggacgcccgg ggcagtgtg agttccagga gggctgtgag  
 2821 ctggtggacg tacgggtgcc cctcttatc cggcctgagg atgacgacga gaagcagctg  
 2881 ctggtggagg ccatcgacgt gccgcaggc actgccacc tcggccgccc cctggtaaac  
 2941 atcaccatca tcaaggagca agccagagac gtggtgtcct ttgagcagcc tgagttctg  
 3001 gtcagccgcg gggaccaggt gggccgcac cctgtcatcc ggcgtgtcct ggacggcggg  
 3061 aagtccagg tctctaccg cacacaggat ggcaccgcg agggcaaccg ggactacatc  
 3121 cccgtggagg gtgagctgct gttcagcct ggggagccct ggaaagagct gcaggtgaag  
 3181 ctctggagc tgcaagaagt tgactcctc ctgcggggcc gccaggtccg ccgtttcac  
 3241 gtccagctca gcaaccctaa gtttggggcc cacctgggac agcccaactc caccaccatc  
 3301 atcatcaggg accagatga actggaccgg agcttcacga gtcagatgtt gtcacacag  
 3361 ccacccctc acggcgacct gggcgcccc cagaaccca atgctaaggc cgctgggtcc  
 3421 aggaagatcc attcaactg gctgcccct tctggcaagc caatgggga cagggtaaag  
 3481 tactggattc agggcgactc cgaatccgaa gccacactgc tcgacagcaa ggtgccctca  
 3541 gtggagctca ccaacctga cccgtattgc gactatgaga tgaaggtgtg cgctacggg

3601 gctcagggcg agggacccta cagctccctg gtgtctgcc gcaccacca ggaagtgcc  
 3661 agcgagccag ggcgtctggc cttaatgtc gtctctcca cggtagacca gctgagctgg  
 3721 gctgagcccg ctgagaccaa cggtagatc acagcctacg aggtctgcta tggcctggc  
 3781 aacgatgaca accgacctat tggggccatg aagaaagtgc tggttgaaa ccctaagaac  
 3841 cggatgtgc ttattgagaa ccttcgggag tcccagccct accgctacac ggtgaaggcg  
 3901 cgcaacgggg cgggctggg gcctgagcgg gaggccatca tcaactggc caccagccc  
 3961 aagaggccca tgtccatccc catcatccct gacatcccta tcgtggacgc ccagagcggg  
 4021 gaggactacg acagcttct tatgtacagc gatgacgtt tacgtctcc atcgggcagc  
 4081 cagaggccca gcgtctcca tgacttggc tgcggctgga agttcgagcc cctgctggg  
 4141 gaggagctgg acctgcggcg cgtcacgtg cggctgccc cggagctcat cccgcgctg  
 4201 tcggccagca gcgggcgctc ctccgacgc gaggcccca cggcccccg gacgacggcg  
 4261 gcgcgggcg gaagggcggc agccgtgcc cgcagtgcga caccgggccc cccgggagag  
 4321 cacctggtga atggccggat ggactttgcc tccccgggca gcaccaactc cctgcacagg  
 4381 atgaccacga ccagtgtgc tgcctatggc acccactga gccacacgt gccccaccg  
 4441 gtgctaagca catctccac cctcacagg gactacaact cactgacccg ctgagaacac  
 4501 tcacactga ccacactgcc cagggactac tcaccctca cctccgtctc ctcccacgac  
 4561 tctgcctga ctgctggtgt gcccacacg cccaccgcc tgggtttctc tgcctgggg  
 4621 cccacatctc tcagagtga ctggcaggag ccgcgggtgc agcgccgct gcagggtac  
 4681 agtgaggagt accagctgt gaacggcggg gagctgcatc ggctcaacat cccaaccct  
 4741 gccagacct cgggtggtgt ggaagacctc ctgcccaacc actctacgt gttccgctg  
 4801 cgggccaga gccaggaagg ctggggccga gacgtgagg gtgtcatcac cattgaatc  
 4861 cagggtcacc cgcagagccc actgtgtccc ctgccaggct ccgccttcac ttgagcact  
 4921 cccagtgcc caggccgct ggtgttact gccctgagcc cagactcgt gcagctgagc  
 4981 tgggagcggc cacggaggcc caatggggat atcgtcggct acctggtgac ctgtgagatg  
 5041 gcccaaggag gagggccagc caccgattc cgggtggatg gagacagccc cgagagccg  
 5101 ctgaccgtgc cgggcctcag cgagaactg ccctacaagt tcaaggtgca ggccaggacc  
 5161 actgagggtc tcgggccaga gcgcgagggc atcatcacca tagagtcca ggatggagga  
 5221 ccttcccg agctgggag ccgtgccggg ctcttcagc accgctgca aagcgagtac  
 5281 agcagcatca ccaccacca caccagcgc accgagccct tcctagtga tgggctgacc  
 5341 ctggggggc agcacctgga ggcaggcggc tcctcacc ggcatgtgac ccaggagttt  
 5401 gtgagccga cactgaccac cagcgaacc ctagaccc acatggacca acagttctc  
 5461 caaactga (Seq. ID. No. 2)

The coding sequence of the alpha 6 chain of human integrin is known from NM\_000210 to be as follows:

1 atggccgccc cggggcagct gtgcttctc tacctgtcgg cggggctct gtcccggctc  
 61 ggcgcagcct tcaacttga cactcgggag gacaactga tccggaaata tggagacccc  
 121 gggagcctct tcggcttctc gctggccatg cactggcaac tgacgcccga ggacaagcgg  
 181 ctgttctcgc tgggggcccc gcgcggagaa gcgcttcac tgacagagc caacagaacg  
 241 ggagggtgt acagctgcga catcaccgcc cgggggcat gcacgcggat cgagtttgat  
 301 aacgatgtc accccacgtc agaaagcaag gaagatcagt ggatgggggt caccgtccag

361 agccaaggct cagggggcaa ggtcgtgaca tgtgctcacc gatatgaaaa aaggcagcat  
 421 gttaatacga agcaggaatc ccgagacatc ttgggcggt gttatgtcct gagtcagaat  
 481 ctgaggattg aagacgatat ggatggggga gattggagct ttgtgatgg gcgattgaga  
 541 ggccatgaga aatttggtc ttgccagcaa ggtgtagcag ctacttttac taaagacttt  
 601 cattacattg tatttgagc cccgggtact tataactgga aagggttgt tcgtgtagag  
 661 caaaagaata acacttttt tgacatgaac atcttgaag atgggcctta tgaagtggg  
 721 ggagagactg agcatgatga aagtctcgtt cctgttctg ctaacagta cttaggtttt  
 781 tctttggact cagggaaagg tattgtttct aaagatgaga tcaactttgt atctggtgct  
 841 cccagagcca atcacagtgg agccgtggtt ttgctgaaga gagacatgaa gtctgcacat  
 901 ctctccctg agcacatatt cgatggagaa ggtctggcct cttcatttgg ctatgatgtg  
 961 gcggtggtgg acctcaaca ggatgggtgg caagatatag ttattggagc cccacagtat  
 1021 ttgatagag atggagaagt tggaggtgca gtgtatgtct acatgaacca gcaaggcaga  
 1081 tggaataatg tgaagccaat tcgtcttaat ggaaccaaag attctatgtt tggcattgca  
 1141 gtaaaaaata ttggagatat taatcaagat ggctaccag atattgcagt tggagctccg  
 1201 tatgatgact tgggaaagg tttatctat catggatctg caaatggaat aaataccaaa  
 1261 ccaacacagg ttctcaagg tatacacct tatttggat attcaattgc tggaaacatg  
 1321 gaccttgatc gaaattccta cctgatgtt gctgttgggt cctctcaga ttcagtaact  
 1381 atttcagat cccggcctgt gattaatatt cagaaaacca tcacagtaac tcctaacaga  
 1441 attgacctc gccagaaaac agcgtgtggg gcgcctagtg ggatagcct ccaggttaaa  
 1501 tcctgtttg aatatactgc taaccccgct ggtataatc ctcaatattc aattgtgggc  
 1561 acactgaag ctgaaaaaga aagaagaaaa tctgggctat cctcaagagt tcagttcga  
 1621 aaccaagggt ctgagcccaa atatactca gaactaact tgaaggagca gaaacagaaa  
 1681 gtgtgcatgg aggaaacct gtggctacag gataatatca gagataaact gcgtccatt  
 1741 cccataactg cctcagtggg gatccaagag ccaagcttc gtaggcgagt gaattcatt  
 1801 ccagaagttc ttcaattct gaattcagat gaaccaaga cagctcatat tgatgtcac  
 1861 ttctaaaag agggatgtgg agacgacaat gtatgaaca gcaacctta actagaatat  
 1921 aaattttgca cccgagaagg aatcaagac aaattttct atttaccat tcaaaaagg  
 1981 gtaccagaac tagttctaaa agatcagaag gatattgctt tagaaataac agtgacaaac  
 2041 agcccttcca acccaaggaa tcccacaaa gatggcgatg acgcccata ggctaaactg  
 2101 attgcaagt ttccagacac tttaacctat tctgcatata gagaactgag ggctttccct  
 2161 gagaaacagt tgagttgtgt tgccaaccag aatggctcgc aagctgactg tgagctcgga  
 2221 aatcctttta aaagaaattc aatgtcact tttatttgg ttttaagtac aactgaagtc  
 2281 accttgaca ccccatatct ggtatatta ctgaagttag aaacaacaag caatcaagat  
 2341 aatttggtc caattacagc taaagcaaaa gtggttattg aactgcttt atcggtctcg  
 2401 ggagttgcta aaccttcca ggtgtatgtt ggaggtacag ttgtggcga gcaagctatg  
 2461 aaatctgaag atgaagtggg aggtttaata gagtatgaat tcagggtaat aaacttaggt  
 2521 aaaccttcta caacctcgg cacagcaacc ttgaacattc agtggccaaa agaaattagc  
 2581 aatgggaaat ggttgcttta ttggtgaaa gtagaatcca aaggattgga aaaggtaact  
 2641 tgtgagccac aaaaggagat aaactccctg aacctaacgg agtctcaca ctcaagaaa  
 2701 aaacgggaaa ttactgaaaa acagatagat gataacagaa aattttctt atttgctgaa  
 2761 agaaaatacc agactctta ctgtagcgtg aacgtgaact gtgtgaacat cagatgccg  
 2821 ctgcgggggc tggacagcaa ggcgtctctt atttgcgtc cgaggttatg gaacagcaca

2881 tttctagagg aatattccaa actgaactac ttggacattc tcatgcgagc cttcattgat  
 2941 gtgactgctg ctgccgaaaa tatcaggctg ccaaatgcag gcactcaggt tcgagtgact  
 3001 gtgtttccct caaagactgt agctcagtat tcgggagtag cttggtggat catcctagt  
 3061 gctattctcg ctgggatctt gatgcttgct ttattagtgt ttatactatg gaagtgtggt  
 3121 ttcttcaaga gaaataagaa agatcattat gatgccacat atcacaaggc tgagatccat  
 3181 gctcagccat ctgataaaga gaggcttact tctgatgcat ag

(Seq. ID. No. 3)

Antisense and RNAi sequence are derivable from these sequences. Antisense oligonucleotides are commonly from 12 to 50 bases in length, more preferably 15-30 bases length. Effective regions for targeting of antisense sequences may be found throughout the target nucleic acid. A preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding Integrin beta 4, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination

codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

RNAi molecules are similarly selected based on the sequence and defined parameters known for the selection of appropriate sequences. RNAi molecules may be single or double stranded, and generally have a length of 19 to 23 bases, although longer and shorter species can be used. A specific RNAi species useful in the method of the invention is based on the mouse sequence of beta-4 cDNA (Genebank Acc. # L04678): nucleotides 113 to 131, counting from the A of the ATG translational start site, having the sequence GAGCTGTACCGAGTGCATC (Seq. ID. No. 4). This molecule, and the corresponding molecule based on the human sequence, and their use form a further aspect of this invention.

In one embodiment of the invention, the therapeutic agent is in combination with other therapy directed toward suppressing the activity of RPTKs known to cooperate with a6b4, including but not limited to ErbB2 (Her2), EGF-R, Met, and Ron. Specific examples of such inhibitors include the Her2 inhibitor trastuzumab (Herceptin™), PPAR gamma ligands as described in US Patent No. 6,291,496, which is incorporated herein by reference

The invention and the evidence that established the efficacy and utility of the invention will now be further described with reference to the following non-limiting examples.

#### Example 1

The MMTV-Neu<sup>Ndl</sup>-YD transgene was introduced into both wild-type and b4-1355T mice of FVB background using the breeding strategy outlined in Fig. 1 (asterick points to Neu mutant. Tumor onset was evaluated by palpation, and mice carrying palpable mammary nodules considered affect. As shown in Fig. 2, the b4 mutant mice lived free of tumors significantly longer than the corresponding control mice. In addition, the b4 mutant mice developed, on average, a smaller number of individual tumors in their

mammary glands. (Fig. 3). These results indicate that  $\alpha 6 \beta 4$  signalling promotes tumorigenesis in this model of breast cancer.

#### Example 2

Tumor growth was evaluated at 6 to 8 weeks after initial detection of tumors. As shown in Fig. 4, the mammary carcinomas of  $\beta 4$  mutant mice grew at a slower rate than those of the control mice, indicating that  $\alpha 6 \beta 4$  signalling promotes tumor growth. Histological analysis indicated that tumors arising in the  $\beta 4$  mutants background were significantly more differentiated (mostly adenocarcinomas) than those arising in the wild-type background (mostly undifferentiated invasive carcinomas). (Fig. 5) Furthermore, immunohistochemistry showed that the  $\beta 4$  mutant retained an apparently intact laminin-containing basement membrane, whereas the wildtype had disrupted the basement membrane and progressed to a frankly invasive stage.

#### Example 3

To begin to examine the mechanism by which  $\alpha 6 \beta 4$  signalling promotes tumor progression, we isolated mammary tumor cell lines from both wild-type and  $\beta 4$  mutant mice. Upon plating on a 2-D matrix, control and  $\beta 4$  mutant tumor cells grew at similar rates. However, when suspended in a Matrigel (a 3-D gel containing basement membrane components) the wild-type tumors proliferated rapidly, producing disorganized aggregates. In contrast, the  $\beta 4$  mutant cells gave rise to small cystic structures resembling normal mammary acini. Taken together, these results provide, for the first time, genetic evidence that  $\alpha 6 \beta 4$  signalling accelerates breast cancer progression by promoting the transition from adenocarcinoma in situ to invasive and metastatic carcinoma and by promoting tumor growth.

#### Example 4

Transgenic mice expressing an SV-40-Tag oncogene from the prostate-specific promoter of Probasin (TRAMP mice) develop prostate cancer with complete penetrance (Gingrich et al., 1992). To examine the role of  $\alpha 6 \beta 4$  signalling in prostate carcinoma progression, we introduced the Probasin-SV-40-Tag transgene in both wild-type and  $\beta 4$ -mutant mice following the breeding strategy outlined in Fig. 6. MRI analysis indicated that tumor onset and growth were delayed in mice carrying the  $\beta 4$  mutation as compared to mice expressing wild-type  $\beta 4$ . (Fig. 7). In addition, the overall survival of the  $\beta 4$  mutant TRAMP mice was observed to be longer than that of the  $\beta 4$ -wild-type TRAMP mice. (Fig. 8).

#### Example 5

Consistent with the results in mammary tumors, histological analysis of prostate tumors from b4-mutant and b4-wild-type TRAMP mice indicated that the tumors arising in the b4-mutant background were considerably more differentiated than those arising in the b4-wild-type background. Anti-Ki67 staining, which marks the nuclei of proliferating cells, revealed that b4-mutant tumors have a significantly reduced proliferative index as compared to the b4-wild-type tumors. Furthermore, b4 was polarized in correspondence of the basement membrane in the tumors of b4-mutant but not b4-wild-type background. Taken together, these findings demonstrate that in the prostate, as in the mammary gland, a6b4 signalling promotes the transition from well differentiated adenocarcinoma with intact basement membrane to invasive carcinoma.

#### Example 6

We have treated with the kinase inhibitor IRESSA, which inhibits both the EGF-R and Neu (de Bono and Rowinsky, 2002), wild-type and  $\beta 4$ -mutant mice carrying MMTV-Neu tumors. The results indicated that IRESSA causes a much larger inhibition of tumor cell proliferation in  $\beta 4$ -mutant mice than it does in wild-type mice (Figure 9). About 80 % of the IRESSA-treated tumors arising in  $\beta 4$ -mutant mice regress, compared to about only 20 % in the control group (Figure 2). This striking result indicates that blockage of  $\alpha 6\beta 4$  increases the effectiveness of cancer therapy with RTK inhibitors, a group which includes all tumor types expressing  $\alpha 6\beta 4$  and carrying amplified or activated versions of Neu, EGF-R, and Met RTKs (Bacus et al., 1994; Longati et al., 2001; Sawyers, 2002).

MMTV-Neu (YD) mice bearing mammary tumors (>0.5 cm in diameter) were treated with Iressa (100 mg/Kg/day) or vehicle (0.1% Tween-80) by gastric gavage for 1 month or 7 days. Tumor sections were stained with anti-Ki-67 Mab, which labels proliferating cells. There is no significant difference in Ki-67 staining between mice treated for 1 month or 7 days. Unpaired, two-tailed t-test showed:  $P < 0.01$  between WT-vehicle and 1355T-vehicle;  $P < 0.001$  between WT-Iressa and 1355T-Iressa;  $P = 0.016$  between 1355T-vehicle and 1355T-Iressa. (Fig. 9)

MMTV-Neu (YD) mice bearing mammary tumors (>0.5cm in diameter) were treated with Iressa (100 mg/Kg/day) by gastric gavage for 24 days. Tumor volumes were measured by caliper. Left panel shows the fold changes in tumor volume from day 0 to day 24, with each line representing one mouse ( $N = 10$  for each group,  $P < 0.01$ ). Right panel of Fig. 10 shows the percentage of mice with regressed tumors in each group.

#### References



The following references are cited herein, and are incorporated herein by reference in their entirety.

- Abdel-Ghany M, Cheng HC, Elble RC, Pauli BU. The breast cancer beta 4 integrin and endothelial human CLCA2 mediate lung metastasis. *J Biol Chem.* 2001 Jul 6;276(27):25438-46.
- Borradori, L., and Sonnenberg, A. (1999). Structure and function of hemidesmosomes: more than simple adhesion complexes. *J. Invest. Dermatol.* 112, 411-418.
- Dans, M., Gagnoux-Palacios, L., Blaikie, P., Klein, S., Mariotti, A., and Giancotti, F. G. (2001). Tyrosine phosphorylation of the b4 integrin cytoplasmic domain mediates Shc signaling to extracellular signal-regulated kinase and antagonizes formation of hemidesmosomes. *J. Biol. Chem.* 276, 1494-1502.
- de Bono, J.S., and E.K. Rowinsky. 2002. The ErbB receptor family: a therapeutic target for cancer. *Trends Mol Med.* 8:519-26.
- Fuchs, E., Dowling, J., Segre, J., Lo, S. H., and Yu, Q. C. (1997). Integrators of epidermal growth and differentiation: distinct functions for b1 and b4 integrins. *Curr. Opin. Genet. Dev.* 7, 672-682.
- Gagnoux-Palacios, L., Dans, M., van't Hof, W., Mariotti, A., Meneguzzi, G., Resh M.D., and Giancotti, F. G. (2003). Integrin  $\alpha 6 \beta 4$  signaling requires compartmentalization in lipid rafts. *J. Cell Biol.* 162, 1189-1196.
- Geiger, B., Bershadsky, A., Pankov, R., and Yamada, K. M. (2001). Transmembrane crosstalk between the extracellular matrix and the cytoskeleton. *Nat. Rev. Mol. Cell Biol.* 2, 793-805.
- Giancotti, F. G., and Tårone, G. (2003). Positional Control of Cell Fate through Joint Integrin/Receptor Protein Tyrosine Kinase Signaling. *Ann. Rev. Cell Dev. Biol.* 19:173-206.
- Giancotti, F. G., and Ruoslahti, E. (1999). Integrin signaling. *Science* 285, 1028-1032.
- Hynes, R. O. (2002). A reevaluation of integrins as regulators of angiogenesis. *Nat. Med.* 8, 918-921.
- Hynes, R. O. (2003). Metastatic potential: generic predisposition of the primary tumor or rare, metastatic variants—or both? *Cell* 113 821-823.

- Mainiero, F., Murgia, C., Wary, K. K., Curatola, A. M., Pepe, A., Blumberg, M., Westwick, J. K., Der, C. J., and Giancotti, F. G. (1997). The coupling of  $\alpha 6 \beta 4$  integrin to Ras-MAP kinase pathways mediated by Shc controls keratinocyte proliferation. *EMBO J.* 16, 2365-2375.
  - Mainiero, F., Pepe, A., Wary, K. K., Spinardi, L., Mohammadi, M., Schlessinger, J., and Giancotti, F. G. (1995). Signal transduction by the  $\alpha 6 \beta 4$  integrin: distinct  $\beta 4$  subunit sites mediate recruitment of Shc/Grb2 and association with the cytoskeleton of hemidesmosomes. *EMBO J.* 14, 4470-4481.
  - Miranti, C. K., and Brugge, J. S. (2002). Sensing the environment: a historical perspective on integrin signal transduction. *Nat. Cell Biol.* 4, E83-90.
  - Murgia, C., Blaikie, P., Kim, N., Dans, M., Petrie, H. T., and Giancotti, F. G. (1998). Cell cycle and adhesion defects in mice carrying a targeted deletion of the integrin  $\beta 4$  cytoplasmic domain. *EMBO J.* 17, 3940-3951.
- Natali PG, Nicotra MR, Botti C, Mottolese M, Bigotti A, Segatto O. Changes in expression of  $\alpha 6 / \beta 4$  integrin heterodimer in primary and metastatic breast cancer. *Br J Cancer.* 1992 Aug;66(2):318-22
- Pellegrini, G., De Luca, M., Orecchia, G., Balzac, F., Cremona, O., Savoia, P., Cancedda, R., and Marchisio, P. C. (1992). Expression, topography, and function of integrin receptors are severely altered in keratinocytes from involved and uninvolved psoriatic skin. *J. Clin. Invest.* 89, 1783-1795.
  - Schlaepfer, D. D., and Hunter, T. (1998). Integrin signalling and tyrosine phosphorylation: just the FAKs? *Trends Cell Biol.* 8, 151-157.
  - Shaw, L. M. (2001). Identification of Insulin Receptor Substrate 1 (IRS-1) and IRS-2 as Signaling Intermediates in the  $\alpha 6 \beta 4$  Integrin-Dependent Activation of PI3-K and Promotion of Invasion. *Mol. Cell Biol.* 21, 5082-5093.
  - Shaw, L. M., Rabinovitz, I., Wang, H. H., Toker, A., and Mercurio, A. M. (1997). Activation of PI3-K by the  $\alpha 6 \beta 4$  integrin promotes carcinoma invasion. *Cell* 91, 949-960.
- Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science.* 1987 Jan 9;235(4785):177-82.

Spinardi, L., Einheber, S., Cullen, T., Milner, T. A., and Giancotti, F. G. (1995). A recombinant tail-less integrin b4 subunit disrupts hemidesmosomes, but does not suppress a6b4-mediated cell adhesion to laminins. *J. Cell Biol.* 129, 473-487.

Tagliabue E, Ghirelli C, Squicciarini P, Aiello P, Colnaghi MI, Menard S. Prognostic value of alpha 6 beta 4 integrin expression in breast carcinomas is affected by laminin production from tumor cells. *Clin Cancer Res.* 1998 Feb;4(2):407-10.

Trusolino, L., Bertotti, A., and Comoglio, P. M. (2001). A signaling adapter function for a6b4 integrin in the control of HGF-dependent invasive growth. *Cell* 107, 643-654.

Wary, K. K., Mariotti, A., Zurzolo, C., and Giancotti, F. G. (1998). A requirement for caveolin-1 and associated kinase Fyn in integrin signaling and anchorage-dependent cell growth. *Cell* 94, 625-634.

Weaver, V. M., Lelievre, S., Lakins, J. N., Chrenek, M. A., Jones, J. C., Giancotti, F., Werb, Z., and Bissell, M. J. (2002). b4 integrin-dependent formation of polarized three-dimensional architecture confers resistance to apoptosis in normal and malignant mammary epithelium. *Cancer Cell* 2, 205-216.

Based on the foregoing, it will be understood that the present invention provides the following advances in the art:

1. A method for inhibition of tumorigenesis in an individual suffering from or at risk for a tumor type that expresses a6b4 integrin, such as thyroid, breast, prostate and cervical cancers, cancer of the upper gastrointestinal tract or squamous carcinoma of the skin, comprising the steps of administering to the individual a therapeutic agent effective to reduce the amount of active a6b4 integrin at least in a portion of the individual where tumorigenesis may occur by targeting the beta 4 portion of the integrin.
2. The method of paragraph 1, wherein the individual is human.
3. The method of paragraph 1 or 2, wherein the therapeutic agent is an antibody.
3. The method of paragraph 1 or 2, wherein the therapeutic agent is an antisense oligonucleotide.
4. The method of paragraph 1 or 2, wherein the therapeutic agent is an RNAi species.

5. The method of any of paragraphs 1 to 4, further comprising the step of administering to the individual an inhibitor of a receptor protein tyrosine kinase such as ErbB2, EGF-R, Met and Ron.
5. Use of an inhibitor of  $\alpha 6 \beta 4$  integrin that targets beta 4 in the preparation of a pharmaceutical composition for inhibition of tumorigenesis.
6. Use of paragraph 5, wherein wherein the therapeutic agent is an antibody.
7. Use of paragraph 5, wherein the therapeutic agent is an antisense oligonucleotide.
8. Use of paragraph 5, wherein the therapeutic agent is an RNAi species.
9. Use of any of paragraphs 5 to 8, wherein the pharmaceutical composition is suitable for human administration.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

# *Introduction of MMTV-Neu\* Transgene in $\beta 4$ 1355T and Wild Type Mice*

$\beta 4^{WT} / \beta 4^{1355T}$  ; MMTV-Neu\* X  $\beta 4^{WT} / \beta 4^{1355T}$  ; MMTV-Neu\*



$\beta 4^{1355T} / \beta 4^{1355T}$  ; MMTV-Neu +  $\beta 4^{WT} / \beta 4^{WT}$  ; MMTV-Neu

Experimental group

Control group

BEST AVAILABLE COPY

Fig. 1

## *Tumor-free Survival on FVB Background*




Fig. 2

# The $\beta$ 4-1355T Mice Develop Fewer Mammary Tumors





Fig 3



# *Reduced Mammary Tumor Growth in $\beta$ 4-1355T Mice*



**Fig. 4**

# *The Mammary Tumors of $\beta$ 4-1355T Mice are more Differentiated*




Fig. 5

# ***Introduction of Probasin-SV40-Tag Transgene in $\beta$ 4<sup>1355T</sup> and Wild Type Mice***

$\beta$ 4<sup>WT</sup> /  $\beta$ 4<sup>1355T</sup> ; TRAMP X  $\beta$ 4<sup>WT</sup> /  $\beta$ 4<sup>1355T</sup> ; TRAMP



$\beta$ 4<sup>1355T</sup> /  $\beta$ 4<sup>1355T</sup> ; TRAMP +  $\beta$ 4<sup>WT</sup> /  $\beta$ 4<sup>WT</sup> ; TRAMP

**Experimental group**

**Control group**

**Fig. 6**

# *MRI Study of Prostate Ca. Progression in $\beta$ 4-1355T Mice*






Fig. 7

***Delayed Progression of Probasin-SV40-TAg-induced  
Prostate Carcinomas in  $\beta$ 4-1355T Mice***



**Fig. 8**

# Loss of $\beta 4$ signaling sensitizes MMTV-Neu tumors to anti-ErbB targeted therapy




Fig. 9

Fig. 10

Fold of Change in Tumor Volume  
After 24-day Iressa Treatment

